


Montana Water Supply Outlook Report March 1, 2007



Picture: Black Bear SNOTEL Site - Winter of 1997

Water Supply Outlook Report

and Federal - State - Private Cooperative Snow Surveys

For more water supply and resource management information, contact:
See Attached List

How forecasts are made

Most of the annual streamflow in the Western United States originates as snowfall that has accumulated high in the mountains during winter and early spring. As the snowpack accumulates, hydrologists estimate the runoff that will occur when it melts. Predictions are based on careful measurements of snow water equivalent at selected index points. Precipitation, temperature, soil moisture and antecedent streamflow data are combined with snowpack data to prepare runoff forecasts. Streamflow forecasts are coordinated by Natural Resources Conservation Service and National Weather Service hydrologists. This report presents a comprehensive picture of water supply conditions for areas dependent upon surface runoff. It includes selected streamflow forecasts, summarized snowpack and precipitation data, reservoir storage data, and narratives describing current conditions.

Snowpack data are obtained by using a combination of manual and automated SNOTEL measurement methods. Manual readings of snow depth and water equivalent are taken at locations called snow courses on a monthly or semi-monthly schedule during the winter. In addition, snow water equivalent, precipitation and temperature are monitored on a daily basis and transmitted via meteor burst telemetry to central data collection facilities. Both monthly and daily data are used to project snowmelt runoff.

Forecast uncertainty originates from two sources: (1) uncertainty of future hydrologic and climatic conditions, and (2) error in the forecasting procedure. To express the uncertainty in the most probable forecast, four additional forecasts are provided. The actual streamflow can be expected to exceed the most probable forecast 50% of the time. Similarly, the actual streamflow volume can be expected to exceed the 90% forecast volume 90% of the time. The same is true for the 70%, 30%, and 10% forecasts. Generally, the 90% and 70% forecasts reflect drier than normal hydrologic and climatic conditions; the 30% and 10% forecasts reflect wetter than normal conditions. As the forecast season progresses, a greater portion of the future hydrologic and climatic uncertainty will become known and the additional forecasts will move closer to the most probable forecast.

The United States Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs and marital or familial status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C., 20250, or call 1-800-245-6340 (voice) or (202) 720-1127 (TDD). USDA is an equal employment opportunity employer.



United States Department of Agriculture
Natural Resources Conservation Service
Bozeman, Montana

(406) 587-6811
10 E Babcock
Federal Bldg Room 443
Bozeman MT 59715

Where to Get More Information

This contact list provides you with the names and phone numbers of individuals who can provide more detailed snow survey information for each county.

Beaverhead County

Victor Hager
683-3800

Dawson County

Linda Peterson-Lohse
365-5566

Hill County

Deana Grabofsky
265-6792

Big Horn County

Evan Van Order
665-3442

Deer Lodge County

Glen Green
846-1703

Jefferson County

Julie Sacks
287-3215

Blaine County

Matthew Crampton
357-2320

Fallon County

Ann Fischer
778-2217

Judith Basin County

Rick Caquelin
566-2311

Broadwater County

Karla Ware
266-3146

Fergus County

Patty Creamer
538-7401

Lake County

David Wolf
676-2841

Carbon County

Gregory Evertz
962-3641

Flathead County

Angel Rosario
752-4242

Lewis and Clark County

Mindy Gauthier
449-5000

Carter County

Wayne Yost
775-6355

Gallatin County

Matt Drechsel
522-4018

Liberty County

Dawn Wickum
759-5778

Cascade County

Mark Delperdang
727-7580

Garfield County

Sue Fitzgerald
557-2232

Lincoln County

Kirk Sullivan
296-2233

Chouteau County

Dan Van Voast
622-5627

Glacier County

Mark Johnson
873-4292

Madison County

Ole Berkram
842-5741

Custer County

Rusty Irion
232-7905

Golden Valley County

Justin Meissner
323-2103

McCone County

Regan Kirchner
485-2744

Daniels County

Valerie Oksendahl
487-5366

Granite County

Mary McDonald
859-3291

Meagher County

Tammy DeCock
547-3633

(over)

Mineral County

Tony Prince
829-3395

Roosevelt County

Robert Bray
787-5232

Wibaux County

Katrina Johnson
796-2211

Missoula County

Tony Prince
829-3395

Rosebud County

Rocky Schwagler
346-7333

Yellowstone County

Valerie Roberson
657-6135

Musselshell County

Justin Meissner
323-2103

Sanders County

Don Feist
826-3701

Park County

Ron Hoagland
222-2899

Sheridan County

Monica Friedrich
765-1801

Petroleum County

Nikki Rife
429-6646

Silver Bow County

Julie Sacks
287-3215

Phillips County

Lorna Philp
654-1334

Stillwater County

Philip Sandoval
322-5359

Pondera County

Stacy Eneboe
278-3922

Sweet Grass County

Chuck Roloff
932-5160

Powder River County

Carol Hilliard
436-2417

Teton County

Thedis Crowe
466-5722

Powell County

Glen Green
846-1703

Toole County

Misty Vermulm
434-5234

Prairie County

Dick Scheetz
635-5381

Treasure County

Danae Pinkerton
342-5510

Ravalli County

Henry Burkwhat
363-5010

Valley County

Denise Wiedenheft
228-4321

Richland County


Jamie Selting
433-2103

Wheatland County

John Oiestad
632-5534

1/8/06

Mountain Snow Water Equivalent



0 50 100 Miles

Current as of
March 1, 2007


NRCS Natural Resources
Conservation Service
<http://www.mt.nrcs.usda.gov>

**NOTE: Data used to generate
this map are PROVISIONAL and
SUBJECT TO CHANGE.**

Percent of Average by Sub-Basin

	Extremely Below Average (Below 51)
	Much Below Average (51 - 70)
	Below Average (71 - 90)
	Near Average (91 - 110)
	Above Average (111 - 130)
	Much Above Average (131 - 150)
	Extremely Above Average (Over 150)

Streamflow Prospects for Montana



Current as of
March 1, 2007

NRCS Natural Resources
Conservation Service
<http://www.mt.nrcs.usda.gov>

NOTE: Data used to generate
this map are **PROVISIONAL** and
SUBJECT TO CHANGE.

STREAMFLOW PROSPECTS

- Extremely Above Average (Over 150)
- Much Above Average (131 - 150)
- Above Average (111 - 130)
- Near Average (91 - 110)
- ◐ Below Average (71 - 90)
- Much Below Average (51 - 70)
- ▲ Extremely Below Average (Below 51)


RIVER INDEX & SWSI VALUES

- 1 Marias above Tiber Reservoir
- 2 Tobacco -0.2
- 3 Kootenai Ft. Steele to Libby Dam 0.5
- 4 Kootenai below Libby Dam 1.7
- 5 Fisher 0.9
- 6 Yaak 1.2
- 7 North FK. Flathead 0.3
- 8 Middle FK. Flathead -0.3
- 9 South FK. Flathead 1.8
- 11 Stillwater/Whitefish 0.6
- 12 Swan -1.1
- 13 Flathead at Polson 0.6
- 14 Mission Valley
- 15 Little Bitterroot
- 16 Clark Fork above Milltown -0.5
- 17 Blackfoot -1.7
- 19 Bitterroot -1.4
- 20 Clark Fork below Bitterroot -1.4
- 21 Clark Fork below Flathead -0.1
- 22 Beaverhead -2
- 23 Ruby -3.6
- 24 Big Hole -0.8
- 25 Boulder (Jefferson) -1
- 26 Jefferson -0.2
- 27 Madison -1.2
- 28 Gallatin -1.6
- 29 Missouri above Canyon Ferry -1.1
- 30 Missouri below Canyon Ferry -1.6
- 31 Smith 0.4
- 32 Sun -0.3
- 33 Teton 0
- 34 Birch/Dupuyer Creeks -3
- 35 Marias
- 36 Musselshell -2.1
- 37 Missouri above Fort Peck
- 38 Missouri below Fort Peck -3.7
- 39 Milk -0.3
- 40 Dearborn near Craig -1.6
- 41 Yellowstone above Livingston 0
- 42 Shields -2.2
- 43 Boulder (Yellowstone) -2.3
- 44 Stillwater -2.2
- 45 Rock/Red Lodge Creeks -3.1
- 46 Clarks Fork -2.1
- 47 Yellowstone above Bighorn -0.9
- 48 Bighorn below Bighorn Lake -1.8
- 49 Little Bighorn -1.1
- 50 Yellowstone below Bighorn -1.3
- 51 Tongue -1.6
- 52 Powder -2
- 53 Upper Judith River -0.7
- 54 Saint Mary River 1.3

Surface Water Supply Index (SWSI) Values

UNITED STATES DEPARTMENT OF AGRICULTURE

NATURAL RESOURCES CONSERVATION SERVICE




Current as of
March 1, 2007

0 45 90 Miles

**NOTE: Data used to generate
this map are PROVISIONAL and
SUBJECT TO CHANGE.**

NRCS Natural Resources
Conservation Service
<http://www.mt.nrcs.usda.gov>



SNOTEL AND SNOW COURSE SUMMARY

MARCH 2007

SNOW COURSE	ELEVATION	DATE	SNOW DEPTH	WATER CONTENT	LAST YEAR	AVERAGE 71-00
ABE LINCOLN	4440	2/26/07	55	16.4	14.3	17.0
ABUNDANCE LAKE	8800	2/25/07	58	15.6	17.3	16.7
ALBRO LAKE SNOTEL	8300	3/01/07	50	13.4	14.9	16.5
AMBROSE	6480	2/27/07	42	9.4	8.3	10.5
ARCH FALLS	7350	2/23/07	37	8.5	9.3	9.0
ASHLEY DIVIDE	4820	2/27/07	22	4.9	5.3	6.2
ASHLEY LAKE	4000	2/27/07	19	4.9	5.2	5.3
BADGER PASS SNOTEL	6900	3/01/07	77	26.8	27.8	29.7
BANFIELD MTN SNOTEL	5600	3/01/07	56	15.2	15.5	17.0
BAREE MIDWAY	4600	2/28/07	84	24.6	30.7	28.7
BAREE TRAIL	3800	2/28/07	44	11.8	12.8	8.2
BARKER LAKES SNOTEL	8250	3/01/07	51	12.2	11.0	11.1
BASIN CREEK SNOTEL	7180	3/01/07	29	6.3	6.6	6.1
BASSOO PEAK	5150	2/23/07	35	9.0	8.7	9.0
BEAGLE SPGS SNOTEL	8850	3/01/07	31	6.2	7.5	7.0
BEAR BASIN	8150	2/25/07	48	11.2	15.0	16.6
BEAVER CREEK SNOTEL	7850	3/01/07	48	11.4	17.7	14.5
BERRY MEADOW	7000	2/28/07	21	3.9	5.4	5.9
BIG SNOWY	7150	2/28/07	73	23.4	19.7	15.3
BISSON CREEK SNOTEL	4920	3/01/07	26	6.7	7.9	8.7
BLACK BEAR SNOTEL	7950	3/01/07	117	28.7	42.4	32.9
BLACK MOUNTAIN	7750	2/27/07	54	13.6	11.7	11.4
BLACK PINE SNOTEL	7100	3/01/07	36	8.6	8.9	10.1
BLACKTAIL	5650	2/27/07	39	10.7	13.1	12.3
BLOODY DICK SNOTEL	7550	3/01/07	41	8.8	10.1	10.6
BLUE LAKE	5900	2/25/07	58	18.5	18.2	21.1
BOTS SOTS	7750	2/23/07	21	4.7	5.2	6.2
BOULDER MTN SNOTEL	7950	3/01/07	63	15.2	20.1	16.6
BOX CANYON SNOTEL	6700	3/01/07	28	7.6	9.0	8.5
BOXELDER CREEK	5100	2/21/07	27	6.6	7.6	6.5
BRACKETT CR SNOTEL	7320	3/01/07	51	10.8	17.5	17.0
BRANHAM LAKES	8850	2/27/07	80	21.5	26.0	23.6
BRUSH CREEK TIMBER	5000	3/01/07	---	6.9E	6.3	7.5
BULL MOUNTAIN	6600	2/23/07	18	3.8	6.2	5.1
BURNT MTN SNOTEL	5880	3/01/07	13	2.9	3.6	5.0
CABIN CREEK	5200	2/25/07	21	5.1	4.4	5.4
CALL ROAD	8050	2/25/07	32	6.7	7.3	9.2
CALVERT CR SNOTEL	6430	3/01/07	31	6.8	9.1	7.5
CAMP SENIA	7890	2/23/07	16	2.9	3.9	4.6
CARROT BASIN SNOTEL	9000	3/01/07	73	17.8	27.3	22.6
CARTER CREEK	7400	2/28/07	16	3.4	4.6	4.2
CHESSMAN RESERVOIR	6200	2/26/07	15	3.6	1.5	3.1
CHICKEN CREEK	4060	2/27/07	52	14.0	17.4	14.4
CLOVER MDW SNOTEL	8800	3/01/07	47	10.2	13.8	14.2
COLE CREEK SNOTEL	7850	3/01/07	34	8.5	8.8	12.0
COLLEY CREEK	6300	2/26/07	25	5.6	7.3	7.0
COMBINATION SNOTEL	5600	3/01/07	20	4.5	4.1	4.5
COPPER BOTTOM SNOTEL	5200	3/01/07	30	6.9	8.3	9.9
COPPER CAMP SNOTEL	6950	3/01/07	104	39.0	41.3	--
COPPER CREEK	5700	2/25/07	25	7.0	9.6	12.5
COPPER MOUNTAIN	7700	2/22/07	31	6.6	9.8	8.9
COTTONWOOD CREEK	6400	2/27/07	29	6.4	5.8	6.0
COYOTE HILL	4200	2/26/07	25	6.0	9.6	9.1
CREVICE MOUNTAIN	8400	2/27/07	38	9.0	9.6	9.0
CRYSTAL LAKE SNOTEL	6050	3/01/07	45	12.9	15.0	9.6
DAD CREEK LAKE	8400	2/25/07	48	9.8	9.9	11.2
DAISY PEAK SNOTEL	7600	3/01/07	27	5.7	7.6	8.7
DALY CREEK SNOTEL	5780	3/01/07	37	9.2	9.1	9.4
DARKHORSE LK. SNOTEL	8700	3/01/07	84	23.0	25.8	26.6

SNOW COURSE	ELEVATION	DATE	SNOW DEPTH	WATER CONTENT	LAST YEAR	AVERAGE 71-00
DAVIS CREEK	5400	2/27/07	66	22.4	20.8	20.7
DEADMAN CR SNOTEL	6450	3/01/07	34	8.0	10.7	8.5
DESERT MOUNTAIN	5600	2/22/07	38	10.0	12.5	12.6
DISCOVERY BASIN	7050	2/23/07	46	9.3	6.3	8.4
DIVIDE SNOTEL	7800	3/01/07	27	5.9	9.0	8.7
DIX HILL	6400	2/25/07	32	8.2	9.5	10.0
DUPUYER CREEK SNOTEL	5750	3/01/07	30	7.4	4.1	9.2
EAGLE CREEK	7000	2/27/07	34	8.2	9.8	--
EAST FORK R.S.	5400	2/27/07	21	4.8	5.9	5.6
EL DORADO MINE	7800	2/24/07	42	9.7	9.2	15.8
ELK HORN SPRINGS	7800	2/25/07	28	5.4	9.3	7.6
ELK PEAK	8000	2/28/07	47	11.5	10.7	12.6
EMERY CREEK SNOTEL	4350	3/01/07	51	13.0	13.6	13.3
FATTY CREEK	5500	3/02/07	62	17.5	22.9	20.4
FISH CREEK	8000	2/26/07	35	8.1	6.6	7.8
FISHER CREEK SNOTEL	9100	3/01/07	97	26.1	35.7	29.6
FIVE-BULL	5700	2/25/07	19	5.3	3.6	5.3
FLATTOP MTN SNOTEL	6300	3/01/07	116	35.9	43.9	39.2
FLEECER RIDGE	7500	2/23/07	30	6.3	8.9	9.2
FOOLHEN	8280	2/25/07	43	9.0	12.3	13.2
FOREST LAKE	6400	2/27/07	29	6.0	8.3	--
FOUR MILE	6900	2/26/07	20	4.0	6.7	6.8
FOURTH OF JULY	3450	2/26/07	36	9.7	7.7	8.4
FREIGHT CREEK	6000	2/25/07	39	10.3	10.4	12.6
FROHNER MDWS SNOTEL	6480	3/01/07	28	6.5	6.2	6.3
GARVER CREEK SNOTEL	4250	3/01/07	41	10.2	8.6	9.1
GOAT MOUNTAIN	7000	2/28/07	33	8.3	8.5	8.7
GRASSHOPPER	7000	2/28/07	21	4.6	4.1	4.7
GRAVE CRK SNOTEL	4300	3/01/07	50	13.6	16.2	14.5
GRIFFIN CR DIVIDE	5150	2/23/07	32	8.3	9.1	9.5
HAND CREEK SNOTEL	5030	3/01/07	40	9.9	10.3	9.9
HAWKINS LAKE SNOTEL	6450	3/01/07	79	25.0	22.9	23.1
HAYMAKER	8050	2/26/07	30	8.2	7.8	--
HEBGEN DAM	6550	2/28/07	29	6.0	8.6	10.8
HELL ROARING DIVIDE	5770	3/01/07	77	24.3	31.1	25.8
HERRIG JUNCTION	4850	2/27/07	67	21.0	27.8	22.2
HOLBROOK	4530	2/28/07	26	8.6	7.6	8.3
HOODOO BASIN SNOTEL	6050	3/01/07	99	31.7	41.0	38.6
INDEPENDENCE	7850	3/01/07	43	11.4	16.3	15.0
INTERGAARD	6450	2/24/07	27	5.3	4.6	6.2
JOHNSON PARK	6450	2/26/07	21	4.6	3.8	5.4
KISHENEHN	3890	2/25/07	39	10.1	8.4	7.3
KRAFT CREEK SNOTEL	4750	3/01/07	32	8.6	12.0	13.6
LAKE CREEK	6100	2/25/07	26	4.9	6.8	7.4
LAKEVIEW CANYON	6930	2/27/07	33	6.2	6.6	8.5
LAKEVIEW RDG. SNOTEL	7400	3/01/07	39	7.9	9.4	9.2
LEMHI RIDGE SNOTEL	8100	3/01/07	37	7.4	9.5	8.7
LICK CREEK SNOTEL	6860	3/01/07	41	9.0	8.1	9.5
LITTLE PARK	7400	3/01/07	---	8.4E	15.9	13.0
LOGAN CREEK	4300	3/01/07	---	6.2E	7.4	6.2
LONE MOUNTAIN SNOTEL	8880	3/01/07	52	12.6	18.2	15.0
LOWER TWIN SNOTEL	7900	3/01/07	45	10.3	15.3	15.1
LUBRECHT SNOTEL	4680	3/01/07	18	4.8	4.4	5.3
LUBRECHT FOREST NO 3	5450	3/01/07	16	3.3	4.3	5.6
LUBRECHT FOREST NO 4	4650	3/01/07	6	1.3	2.4	2.7
LUBRECHT FOREST NO 6	4040	3/01/07	8	1.9	3.3	3.2
LUBRECHT HYDROPLOT	4200	3/01/07	20	4.2	5.0	5.1
MADISON PLT SNOTEL	7750	3/01/07	71	16.2	22.7	20.9
MANY GLACIER SNOTEL	4900	3/01/07	42	10.9	10.2	14.4
MARIAS PASS	5250	2/27/07	47	14.2	14.8	14.9
MIDDLE MILL CREEK	7850	2/27/07	47	10.0	14.1	13.3
MILL CREEK	7500	2/26/07	45	11.2	11.4	10.2
MINERAL CREEK	4000	2/27/07	47	13.6	17.3	15.8
MONUMENT PK SNOTEL	8850	3/01/07	59	15.3	20.7	17.2

SNOW COURSE	ELEVATION	DATE	SNOW DEPTH	WATER CONTENT	LAST YEAR	AVERAGE 71-00
MOSS PEAK SNOTEL	6780	3/01/07	96	30.4	34.7	31.3
MOULTON RESERVOIR	6850	2/26/07	27	5.2	7.6	6.2
MT LOCKHART SNOTEL	6400	3/01/07	54	17.0	16.6	17.6
MULE CREEK SNOTEL	8300	3/01/07	45	10.2	12.7	13.1
N.E. ENTRANCE SNOTEL	7350	3/01/07	33	7.8	9.0	9.7
NEVADA RIDGE SNOTEL	7020	3/01/07	40	9.9	13.2	13.2
NEW WORLD	6900	2/23/07	40	10.2	12.4	11.1
NEWTON MOUNTAIN	5600	2/26/07	73	26.5	28.2	28.8
NEZ PERCE CMP SNOTEL	5650	3/01/07	42	10.3	14.0	12.7
NEZ PERCE CREEK	6600	2/22/07	19	3.5	6.1	5.6
NEZ PERCE PASS	6570	2/28/07	41	10.2	14.2	15.7
N.F. ELK CR SNOTEL	6250	3/01/07	37	9.0	9.4	10.2
NF JOCKO SNOTEL	6330	3/01/07	88	28.9	39.7	38.5
NOISY BASIN SNOTEL	6040	3/01/07	95	29.6	41.7	33.8
NOTCH	8500	2/25/07	53	11.8	8.6	12.4
OPHIR PARK	7150	2/25/07	36	9.4	13.0	14.1
PETERSON MDW SNOTEL	7200	3/01/07	38	8.2	7.2	7.8
PICKET PIN LOWER	6200	2/28/07	10	2.1	1.3	--
PICKET PIN MIDDLE	7250	2/28/07	19	3.7	7.3	--
PICKET PIN UPPER	8100	2/28/07	41	11.8	15.5	--
PICKFOOT CRK SNOTEL	6650	3/01/07	36	7.9	11.4	9.9
PIKE CREEK SNOTEL	5930	3/01/07	61	19.6	23.6	22.8
PIPESTONE PASS	7200	2/22/07	16	3.2	4.2	4.1
PLACER BASIN SNOTEL	8830	3/01/07	48	12.1	16.0	14.2
POORMAN CR SNOTEL	5100	3/01/07	97	31.6	37.5	28.2
PORCUPINE SNOTEL	6500	3/01/07	21	4.5	7.1	5.8
POTOMAGETON PARK	7150	2/28/07	42	9.8	14.2	12.1
RED TOP	5260	2/28/07	69	22.5	22.0	23.8
REVAIS CREEK	4800	2/27/07	6	1.0	.2	3.1
ROCK CREEK	5600	2/28/07	36	10.0	12.2	8.4
ROCK CREEK MEADOW	8160	2/26/07	54	13.6	21.3	16.6
ROCKER PEAK SNOTEL	8000	3/01/07	45	10.3	12.1	11.2
ROCKY BOY SNOTEL	4700	3/01/07	23	5.0	4.4	4.3
SACAJAWEA	6550	2/26/07	28	6.6	10.3	11.2
SACAJAWEA SNOTEL	6550	3/01/07	41	8.9	12.5	12.0
SADDLE MTN SNOTEL	7900	3/01/07	71	17.1	23.9	21.8
S.F. SHIELDS SNOTEL	8100	3/01/07	48	10.6	16.1	13.6
SHORT CREEK SNOTEL	7000	3/01/07	18	4.0	5.2	4.9
SHOWER FALLS SNOTEL	8100	3/01/07	67	16.2	19.0	17.4
SKALKAHO SNOTEL	7260	3/01/07	62	17.6	21.3	20.2
SLEEPING WOMAN SNTL	6150	3/01/07	46	12.5	14.2	14.0
SLIDE ROCK MOUNTAIN	7100	2/25/07	37	9.4	11.3	12.6
SMUGGLER MINE	6960	2/27/07	32	5.6	8.6	8.1
SPOTTED BEAR MTN.	7000	2/24/07	34	9.0	11.0	12.7
SPUR PARK SNOTEL	8100	3/01/07	67	15.9	19.0	17.7
STAHL PEAK SNOTEL	6030	3/01/07	87	27.5	34.2	29.9
STEMPLE PASS	6600	2/26/07	29	6.8	8.2	8.3
STORM LAKE	7780	2/23/07	52	10.0	10.8	10.2
STRYKER BASIN	6180	2/27/07	76	25.8	35.5	26.9
STUART MOUNTAIN SNTL	7400	3/01/07	87	26.7	30.5	28.9
TAYLOR ROAD	4080	2/21/07	17	3.9	3.3	2.7
TEN MILE LOWER	6600	2/28/07	27	6.0	6.4	5.9
TEN MILE MIDDLE	6800	2/28/07	32	7.5	8.7	8.9
TEPEE CREEK SNOTEL	8000	3/01/07	51	10.8	13.8	10.7
TIMBERLINE CREEK	8850	2/23/07	37	9.6	8.0	11.0
TIZER BASIN SNOTEL	6840	3/01/07	36	9.5	8.2	7.9
TRAIL CREEK	7090	2/25/07	27	5.1	7.9	7.0
TRINKUS LAKE	6100	2/24/07	93	32.1	36.7	36.4
TRUMAN CREEK	4060	2/28/07	22	5.3	4.3	4.4
TV MOUNTAIN	6800	3/02/07	52	14.4	17.6	15.0
TWELVEMILE SNOTEL	5600	3/01/07	49	13.7	19.2	16.0
TWENTY-ONE MILE	7150	2/27/07	51	10.8	17.3	14.3
TWIN CREEKS	3580	2/24/07	26	7.5	9.7	10.2
TWIN LAKES SNOTEL	6400	3/01/07	98	33.0	42.5	34.7

SNOW COURSE	ELEVATION	DATE	SNOW DEPTH	WATER CONTENT	LAST YEAR	AVERAGE 71-00
UPPER HOLLAND LAKE	6200	2/24/07	77	23.2	27.4	30.0
WALDRON SNOTEL	5600	3/01/07	38	9.7	9.9	9.7
WARM SPRINGS SNOTEL	7800	3/01/07	71	18.1	17.6	17.0
WEASEL DIVIDE	5450	2/28/07	82	27.2	32.2	28.7
WEST YELL'ST SNOTEL	6700	3/01/07	40	8.3	12.2	11.2
WHISKEY CREEK SNOTEL	6800	3/01/07	55	11.1	13.6	14.4
WHITE MILL SNOTEL	8700	3/01/07	63	16.4	22.8	20.3
WHITE PINE RIDGE	8850	2/25/07	25	4.5	4.7	4.7
WILLOW CREEK	6500	2/26/07	19	3.6	5.8	6.3
WOOD CREEK SNOTEL	5960	3/01/07	31	7.6	9.5	8.2
WRONG CREEK	5700	2/27/07	41	8.9	10.2	11.0
WRONG RIDGE	6800	3/01/07	---	13.0E	14.8	15.4

Montana Water Supply Outlook Report as of March 1, 2007

February has been the best snowfall month so far this winter, increasing 7 percent from February 1. February accumulated snow water equivalent, as a percent of the seasonal peak, in the Columbia River Basin was 118 percent of average, St. Mary was 100 percent of average, Missouri was 131 percent of average, and Yellowstone was 113 percent of average. February mountain precipitation was 105 percent of average and water year precipitation was 101 percent of average. February mountain precipitation in the Columbia was 93 percent of average, Missouri was 118 percent of average, St. Mary was 73 percent of average, and Yellowstone was 111 percent of average.

Snowpack

March 1 is more than three quarters of the way through the main snowfall period and there are normally about four to six weeks remaining until the seasonal snowpack peak is reached. On March 1, accumulated snow water equivalent west of the Divide is about 82 percent and east of the Divide is about 75 percent, leaving about 20 to 25 percent of the seasonal snowfall period remaining. State-wide, mountain snow water content was 87 percent of average and 84 of last year at this time. West of the Continental Divide, snowpack was 90 percent of average and 87 percent of last year and east of the Continental Divide, snowpack was 84 percent of average and 85 percent of last year.

March is still a critical snowfall month. Having an average to above average March snowfall will build on a very good February and bring snowpacks closer to average.

RIVER BASIN	% OF AVERAGE	% OF LAST YEAR	FEBRUARY % CHANGE
COLUMBIA	90	87	+5
KOOTENAI, MONTANA	99	93	0
KOOTENAY, CANADA	109	122	-3
FLATHEAD, MONTANA	89	84	+5
FLATHEAD, CANADA	102	95	+4
UPPER CLARK FORK	85	89	+8
BITTERROOT	85	79	+5
LOWER CLARK FORK	93	86	+5
MISSOURI	87	84	+16
MISSOURI HEADWATERS	82	77	+11
JEFFERSON	83	81	+9
MADISON	79	74	+11
GALLATIN	80	72	+16
MISSOURI MAINSTEM	98	97	+15
HEADWATERS MAINSTEM	93	89	+11
SMITH-JUDITH-MUSSELSHELL ..	102	88	+23
SUN-TETON-MARIAS	90	97	0
MILK	136	155	---
BEARPAW MOUNTAINS	115	101	+42
CYPRESS HILLS, CANADA ...	152	240	+48
ST. MARY	87	85	+1
ST. MARY & MILK	102	105	+20
YELLOWSTONE	79	85	+9
UPPER YELLOWSTONE	81	79	+11
LOWER YELLOWSTONE	77	92	+7
STATE-WIDE	87	84	+7

Precipitation

February mountain and valley precipitation across the state was 117 percent of average and 118 percent of last year, while the water year precipitation was 105 percent of average and 93 percent of last year. West of the Continental Divide, February mountain and valley precipitation was 102 percent of average and 101 percent of last year and the water year precipitation was 107 percent of average and 97 percent of last year. East of the Divide, February mountain and valley precipitation was 131 percent of average and 135 percent of last year and the water year precipitation was 103 percent of average and 90 percent of last year.

RIVER BASIN	FEBRUARY % OF AVERAGE	WATER YEAR % OF AVERAGE
COLUMBIA	102	107
KOOTENAI	85	115
FLATHEAD	105	105
UPPER CLARK FORK	112	104
BITTERROOT	96	103
LOWER CLARK FORK	95	107
MISSOURI	132	104
JEFFERSON	124	104
MADISON	122	94
GALLATIN	142	95
MISSOURI MAINSTEM	160	106
SMITH-JUDITH-MUSSELSHELL	151	114
SUN-TETON-MARIAS	102	103
MILK	198	127
ST. MARY	83	108
YELLOWSTONE	121	95
UPPER YELLOWSTONE	121	100
LOWER YELLOWSTONE	127	91
STATE-WIDE	117	105

Reservoirs

Major reservoir storages state-wide were 82 percent of average and 95 percent of last year. Reservoir storage west of the Continental Divide was 137 percent of average and 91 percent of last year. East of the Continental Divide, reservoir storages were 67 percent of average and 98 percent of last year.

RIVER BASIN	% OF AVERAGE	% OF LAST YEAR
COLUMBIA	137	91
KOOTENAI	155	81
FLATHEAD	127	101
UPPER CLARK FORK	98	90
BITTERROOT	168	187
LOWER CLARK FORK	109	105
MISSOURI	66	98
JEFFERSON	90	119
MADISON	105	100
GALLATIN	153	103
MISSOURI MAINSTEM	63	98
SMITH-JUDITH-MUSSELSHELL	64	94
SUN-TETON-MARIAS	96	95
MILK	79	80
ST. MARY	139	92
YELLOWSTONE	96	94
UPPER YELLOWSTONE	98	85
LOWER YELLOWSTONE	96	94
STATE-WIDE	82	95

Streamflow

Streamflows across Montana are forecast to average 73 to 87 percent of normal. West of the Continental Divide, streamflows are forecast to average 84 to 94 percent of normal and east of the Continental Divide, streamflows are forecast to average 66 to 83 percent of normal.

Below are averaged streamflow forecasts, by river basin, for the period April 1 through July 31. THESE FORECASTS ASSUME NEAR NORMAL SPRING CONDITIONS AND DO NOT ACCOUNT FOR WELL BELOW AVERAGE (70% or less) OR ABOVE AVERAGE (110% or more) SNOWMELT OR SPRING RAIN. Specific forecast probabilities are available in each individual River Basin Report on the Montana NRCS web page.

RIVER BASIN	April-July THIS YEAR		April-July LAST YEAR	
	% OF AVERAGE		% OF AVERAGE	
COLUMBIA	84 to	94	89 to	100
KOOTENAI	90 to	98	90 to	99
FLATHEAD	87 to	95	95 to	104
UPPER CLARK FORK	80 to	94	78 to	94
BITTERROOT	80 to	88	103 to	112
LOWER CLARK FORK	84 to	93	89 to	98
MISSOURI	64 to	83	78 to	98
JEFFERSON	56 to	76	70 to	91
MADISON	77 to	85	103 to	111
GALLATIN	74 to	84	99 to	110
MISSOURI MAINSTEM	67 to	80	87 to	102
SMITH-JUDITH-MUSSELSHELL .	48 to	73	99 to	119
SUN-TETON-MARIAS	76 to	95	68 to	86
MILK	67 to	101	46 to	86
ST. MARY	92 to	98	85 to	92
YELLOWSTONE	69 to	81	88 to	100
UPPER YELLOWSTONE	76 to	87	98 to	108
LOWER YELLOWSTONE	60 to	74	75 to	89
STATE-WIDE	73 to	87	84 to	99

NOTE: The *APRIL-JULY LAST YEAR % OF AVERAGE* column above is what was forecast March 1 last year, NOT what actually occurred.

Surface Water Supply Index


The Surface Water Supply Index (SWSI) is a measure of available surface water availability for the spring and summer months. Water users that rely on mountain precipitation can use the index to evaluate seasonal surface water supplies. The SWSI accounts for mountain snowpack, mountain precipitation, streamflow, reservoir storage, and soil moisture.

SWSI RATING	SURFACE WATER CONDITION
+3.0 to +4.0	Extremely Wet
+2.0 to +2.9	Moderately Wet
+1.0 to +1.9	Slightly Wet
-0.9 to +0.9	Near Average
-1.0 to -1.9	Slightly Dry
-2.0 to -2.9	Moderately Dry
-3.0 to -4.0	Extremely Dry

This year SWSI	Last year SWSI	Basin
-0.2	+0.7	Tobacco River
+0.5	0.0	Kootenai Ft. Steele to Libby Dam
+1.7	+3.0	Kootenai River below Libby Dam
+0.9	+1.1	Fisher River
+1.2	+0.9	Yaak River
+0.3	+0.7	North Fork Flathead River
-0.3	+0.3	Middle FORK Flathead River
+1.8	+3.1	South Fork Flathead River
+1.2	+1.3	Flathead River at Columbia Falls
+0.6	+1.4	Stillwater/Whitefish Rivers
-1.1	+0.9	Swan River
+0.6	+0.9	Flathead River at Polson
----	+0.3	Mission Valley
----	+0.7	Little Bitterroot River
-0.5	+0.3	Clark Fork River above Milltown
-1.7	-1.7	Blackfoot River
-1.4	+1.8	Bitterroot River
-1.4	+0.4	Clark Fork River below Bitterroot River
-0.1	+0.7	Clark Fork River below Flathead River
-2.0	-1.4	Beaverhead River
-3.6	+0.6	Ruby River
-0.8	+0.5	Big Hole River
-1.0	+0.8	Boulder River (Jefferson)
-0.2	+1.4	Jefferson River
-1.2	+0.9	Madison River
-1.6	+1.9	Gallatin River
-1.1	+0.3	Missouri River above Canyon Ferry
-1.6	-0.2	Missouri River below Canyon Ferry
+0.4	+1.6	Smith River
-0.3	-1.6	Sun River
0.0	-1.5	Teton River
-3.0	-2.8	Birch/Dupuyer Creeks
-0.7	+1.8	Upper Judith River
----	-1.6	Marias River above Tiber
----	+0.1	Marias River below Tiber
-2.1	+1.6	Musselshell River
----	-0.9	Missouri River above Ft. Peck
-3.7	-3.4	Missouri River below Ft. Peck
+1.3	+0.4	St. Mary River
-0.3	-1.8	Milk River
-1.6	-1.1	Dearborn River near Craig
0.0	+1.3	Yellowstone River above Livingston
-2.2	+2.3	Shields River
-2.3	+2.5	Boulder River (Yellowstone)
-2.2	+1.3	Stillwater River
-3.1	-1.3	Rock/Red Lodge Creeks
-2.1	+0.7	Clarks Fork River
-0.9	+1.3	Yellowstone River above Bighorn River
-1.8	-0.5	Bighorn River below Bighorn Lake
-1.1	-0.2	Little Bighorn River
-1.3	+0.5	Yellowstone River below Bighorn River
-1.6	-0.5	Tongue River
-2.0	-0.2	Powder River


Kootenai River Basin in Montana

Snowpack conditions in the Kootenai River Basin were average and is the same percent of average as on February 1. Snow water content was 99 percent of average and 1 percent of last year. Snow water content in the Kootenay in Canada was 109 percent of average and 122 percent of last year and decreased 3 percent from February 1. February accumulated snow water equivalent, as a percent of the seasonal peak, was 112 percent of average.



January maximum swe was established in 1997 and minimum was in 1977; February maximum swe was in 1997 and minimum swe was in 1977; March maximum swe was in 1972 and minimum swe was in 1977; April maximum swe was in 1974 and minimum swe was in 2001 and 2005; May maximum swe was in 1974 and minimum swe was in 1977; and June maximum swe was in 1974 and minimum swe was in 1992. Average is for the period 1971 through 2000.

Mountain precipitation during February was 87 percent of average and 86 percent of last year. Valley precipitation during February was 66 percent of average and 83 percent of last year. Mountain and valley water year precipitation, beginning October 1, 2006, was 115 percent of average and 104 percent of last year.



Lake Koocanusa storage was 155 percent of average and 81 percent of last year.

Assuming average precipitation March through July, April through July streamflows are forecast to average 90 percent to 98 percent.

Surface Water Supply Index (SWSI) was -0.2 in the Tobacco River; +0.5 in the Kootenai Ft. Steele to Libby Dam; +1.7 in the Kootenai River below Libby Dam; +0.9 in the Fisher River; and +1.2 in the Yaak River.

KOOTENAI RIVER BASIN in Montana
Streamflow Forecasts - March 1, 2007

		<<----- Drier ----- Future Conditions ----- Wetter ----->>						
Forecast Point	Forecast Period	Chance Of Exceeding *						30-Yr Avg (1000AF)
		90% (1000AF)	70% (1000AF)	50% (1000AF)	(% AVG.)	30% (1000AF)	10% (1000AF)	
TOBACCO RIVER nr Eureka	APR-JUL	87	108	122	90	136	157	136
	APR-SEP	95	119	135	90	151	175	150
LIBBY RES Inflow (1,2)	APR-JUL	4720	5290	5700	101	6120	6770	5640
	APR-SEP	5660	6270	6700	101	7150	7830	6640
FISHER RIVER nr Libby	APR-JUL	157	193	220	96	250	295	230
	APR-SEP	170	210	235	96	265	310	245
YAAK RIVER nr Troy	APR-JUL	385	440	480	103	520	585	465
	APR-SEP	400	455	495	101	535	605	490
KOOTENAI at Leonia (1,2)	APR-JUL	6030	6760	7090	101	7420	8150	7040
	APR-SEP	7690	7980	8180	101	8380	8670	8120

KOOTENAI RIVER BASIN in Montana
Reservoir Storage (1000 AF) - End of February

Reservoir	Usable Capacity	*** Usable Storage ***		
		This Year	Last Year	Avg
LAKE KOOCANUSA	5748.0	3025.0	3751.0	1950.0

KOOTENAI RIVER BASIN in Montana
Watershed Snowpack Analysis - March 1, 2007

Watershed	Number of Data Sites	This Year as % of	
		Last Yr	Average
KOOTENAY in CANADA	15	125	105
KOOTENAI MAINTSTEM	3	89	100
TOBACCO	3	83	93
FISHER	4	89	98
YAAK	7	107	101
KOOTENAI in MONTANA	17	93	99
KOOTENAI ab BONNERS FERRY	32	102	101


* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average is computed for the 1971-2000 base period.

- (1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
- (2) - The value is natural volume - actual volume may be affected by upstream water management.
- (3) - Median value used in place of average. The value listed under 30% is actually a 25% exceedance level. The value listed under 70% is actually a 75% exceedance level. Forecast issued in cooperation with Alberta Environment.


Flathead River Basin

Snowpack conditions in the Flathead River Basin were below average and increased 5 percent from February 1. Snow water content was 89 percent of average and 84 percent of last year. Snow water content in the North Fork Flathead River in Canada was 102 percent of average and 95 percent of last year and increased 4 percent from February 1. February accumulated snow water equivalent, as a percent of the seasonal peak, was 112 percent of average.



January maximum swe was established in 1997 and minimum was in 2001; February maximum swe was in 1972 and minimum was in 2001; March maximum swe was in 1972 and minimum was in 2005; April maximum swe was in 1972 and minimum was in 2005; May maximum swe was in 1972 and minimum was in 1992; and June maximum swe was in 1974 and minimum was in 1992. Average is for the period 1971 through 2000.

Mountain precipitation during February was 103 percent of average and 90 percent of last year. Valley precipitation during February was 143 percent of average and 153 percent of last year. Mountain and valley water year precipitation, beginning October 1, 2006, was 105 percent of average and 92 percent of last year.



Combined Camas reservoir storage was not available; Lower Jocko Lake was not available; combined Mission Valley reservoir storage was not available; Hungry Horse storage was 137 percent of average and 102 percent of last year; and Flathead Lake storage was 102 percent of average and 98 percent of last year.

Assuming average precipitation March through July, April through July streamflows are forecast to average 87 percent to 95 percent.

Surface Water Supply Index (SWSI) was +0.3 in the North Fork Flathead River; -0.3 in the Middle Fork Flathead River; +1.8 in the South Fork Flathead River; +1.2 in the Flathead River at Columbia Falls; +0.6 in the Stillwater/Whitefish Rivers; -1.1 in the Swan River; +0.6 in the Flathead River at Polson; ---- in the Mission Valley; and ---- in the Little Bitterroot River.

FLATHEAD RIVER BASIN
Streamflow Forecasts - March 1, 2007

		<<----- Drier ----- Future Conditions ----- Wetter ----->>						
Forecast Point	Forecast Period	Chance Of Exceeding *						30-Yr Avg. (1000AF)
		90% (1000AF)	70% (1000AF)	50% (1000AF)	(% AVG.)	30% (1000AF)	10% (1000AF)	
NF FLATHEAD nr Columbia Falls	APR-JUL	1370	1500	1590	98	1680	1810	1620
	APR-SEP	1530	1670	1770	98	1870	2010	1800
MF FLATHEAD nr West Glacier	APR-JUL	1270	1420	1520	96	1630	1790	1590
	APR-SEP	1390	1550	1660	95	1770	1950	1740
HUNGRY HORSE Reservoir Inflow (1,2)	APR-JUL	1560	1720	1840	92	1960	2140	2000
	APR-SEP	1660	1830	1950	92	2070	2260	2120
FLATHEAD at Columbia Falls (2)	APR-JUL	4330	4770	5090	95	5420	5910	5350
	APR-SEP	4720	5190	5530	95	5880	6410	5820
STILLWATER nr Whitefish	APR-JUL	137	165	185	96	205	240	192
	APR-SEP	151	182	205	98	230	270	210
WHITEFISH nr Kalispell	APR-JUL	78	91	100	99	110	125	101
	APR-SEP	87	102	112	98	123	140	114
ASHLEY CREEK nr Marion (2)	APR-JUL	3.5	5.4	6.8	94	8.4	11.1	7.2
SWAN RIVER nr Bigfork	APR-JUL	410	460	495	88	530	590	565
	APR-SEP	465	520	560	87	600	665	645
Flathead Lake inflow (1,2)	APR-JUL	5140	5660	6030	98	6410	7000	6180
	APR-SEP	5570	6140	6540	98	6960	7590	6700

FLATHEAD RIVER BASIN
Reservoir Storage (1000 AF) - End of February

Reservoir	Usable Capacity	*** Usable Storage ***		
		This Year	Last Year	Avg
CAMAS (4)		NO REPORT		
LOWER JOCKO LAKE		NO REPORT		
MISSION VALLEY (8)		NO REPORT		
HUNGRY HORSE	3451.0	2799.0	2748.0	2047.6
FLATHEAD LAKE	1791.0	822.1	840.2	802.7

FLATHEAD RIVER BASIN
Watershed Snowpack Analysis - March 1, 2007

Watershed	Number of Data Sites	This Year as % of	
		Last Yr	Average
NF FLATHEAD in CANADA	1	96	93
NF FLATHEAD in MONTANA	9	83	93
MIDDLE FORK FLATHEAD	6	88	90
SOUTH FORK FLATHEAD	7	83	85
STILLWATER-WHITEFISH	10	81	94
SWAN	7	79	83
MISSION VALLEY	4	83	90
LITTLE BITTERROOT-ASHLEY	6	94	92
JOCKO	5	82	84
FLATHEAD in MONTANA	40	84	89
FLATHEAD RIVER BASIN	41	84	89

* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.


The average is computed for the 1971-2000 base period.

- (1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
- (2) - The value is natural volume - actual volume may be affected by upstream water management.
- (3) - Median value used in place of average. The value listed under 30% is actually a 25% exceedance level. The value listed under 70% is actually a 75% exceedance level. Forecast issued in cooperation with Alberta Environment.

Upper Clark Fork River Basin

Snowpack conditions in the Upper Clark Fork River Basin were below average and increased 8 percent from February 1. Snow water content was 85 percent of average and 89 percent of last year. February accumulated snow water equivalent, as a percent of the seasonal peak, was 123 percent of average. Snowpack in the Blackfoot is ninth lowest of record, but has improved from fifth lowest of record on February 1.


Upper Clark Fork Snow Water Equivalent



January maximum swe was established in 1997 and minimum swe was in 1977; February maximum was in 1972 and minimum swe was in 1977; March maximum swe was in 1972 and minimum swe was in 2005; April maximum swe was in 1972 and minimum was in 2005; May maximum swe was in 1972 and minimum swe was in 1977; and June maximum swe was in 1975 and minimum swe was in 1987. Average is for the period 1971 through 2000.

Mountain precipitation during February was 109 percent of average and 125 percent of last year. Valley precipitation during February was 149 percent of average and 203 percent of last year. Mountain and valley water year precipitation, beginning October 1, 2006, was 104 percent of average and 102 percent of last year.

Upper Clark Fork Precipitation



East Fork Rock Creek was 68 percent of average and 72 percent of last year; Georgetown Lake was 108 percent of average and 97 percent of last year; Lower Willow Creek storage was not available; and Nevada Creek storage was 100 percent of average and 84 percent of last year.

Assuming average precipitation March through July, April through July streamflows are forecast to average 80 percent to 94 percent.

Surface Water Supply Index (SWSI) was -0.5 in the Clark Fork River above Milltown; -1.7 in the Blackfoot River; and -1.4 in the Clark Fork River above Missoula.

UPPER CLARK FORK RIVER BASIN
Streamflow Forecasts - March 1, 2007

		<<----- Drier ----- Future Conditions ----- Wetter ----->>						
Forecast Point	Forecast Period	Chance Of Exceeding *						30-Yr Avg. (1000AF)
		90% (1000AF)	70% (1000AF)	50% (1000AF)	(% AVG.)	30% (1000AF)	10% (1000AF)	
=====								
LITTLE BLACKFOOT nr Garrison	APR-JUL	41	62	78	103	96	126	76
	APR-SEP	46	69	86	102	105	138	84
FLINT CREEK nr Southern Cross	APR-JUL	8.7	11.6	13.8	101	16.2	20	13.7
	APR-SEP	9.8	13.4	16.2	100	19.2	24	16.2
FLINT CREEK blw Boulder Ck	APR-JUL	33	45	54	96	64	80	56
	APR-SEP	45	58	69	97	80	99	71
LOWER WILLOW CK RES Inflow	APR-MAY	4.2	6.0	7.3	89	8.8	11.2	8.2
	APR-JUL	5.6	8.4	10.7	86	13.2	17.4	12.5
MF ROCK CREEK nr Philipsburg	APR-JUL	47	56	62	97	69	79	64
	APR-SEP	52	62	69	96	76	88	72
ROCK CREEK nr Clinton	APR-JUL	181	225	260	96	295	355	270
	APR-SEP	210	260	295	97	335	395	305
CLARK FORK abv Milltown	APR-JUL	370	495	595	98	700	875	605
	APR-SEP	445	585	695	99	810	1000	705
NEVADA CREEK nr Helmville	APR-MAY	4.5	6.6	8.3	81	10.2	13.3	10.3
	APR-JUL	7.4	11.2	14.2	82	17.6	23	17.3
BLACKFOOT RIVER nr Bonner	APR-JUL	520	615	685	85	755	870	805
	APR-SEP	585	685	760	85	835	960	890
CLARK FORK abv Missoula	APR-JUL	925	1150	1300	92	1450	1670	1410
	APR-SEP	1060	1310	1470	92	1630	1880	1600

UPPER CLARK FORK RIVER BASIN
Reservoir Storage (1000 AF) - End of February

Reservoir	Usable Capacity	*** Usable Storage ***		
		This Year	Last Year	Avg
EAST FORK ROCK CREEK	15.6	6.4	8.9	9.4
GEORGETOWN LAKE	31.0	28.8	29.7	26.6
LOWER WILLOW CREEK		NO REPORT		
NEVADA CREEK	12.6	5.2	6.2	5.2

UPPER CLARK FORK RIVER BASIN
Watershed Snowpack Analysis - March 1, 2007

Watershed	Number of Data Sites	This Year as % of	
		Last Yr	Average
CLARK FORK ab FLINT CREEK	14	91	89
FLINT CREEK	6	110	97
ROCK CREEK	5	93	89
CLARK FORK ab BLACKFOOT	22	95	90
BLACKFOOT	15	82	76
UPPER CLARK FORK BASIN	35	89	85

* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.


The average is computed for the 1971-2000 base period.

- (1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
- (2) - The value is natural volume - actual volume may be affected by upstream water management.
- (3) - Median value used in place of average. The value listed under 30% is actually a 25% exceedance level. The value listed under 70% is actually a 75% exceedance level. Forecast issued in cooperation with Alberta Environment.

Bitterroot River Basin

Snowpack conditions in the Bitterroot River Basin were below average and increased 5 percent from February 1. Snow water content was 85 percent of average and 79 percent of last year. February accumulated snow water equivalent, as a percent of the seasonal peak, was 111 percent of average.


Bitterroot Snow Water Equivalent



January maximum swe was established in 1997 and minimum swe in 1977; February maximum swe was in 1972 and minimum was in 1977; March maximum swe was in 1972 and minimum swe was in 1977 and 2005; April maximum swe was in 1972 and minimum swe was in 2005; May maximum swe was in 1972 and minimum swe was in 1987; and June maximum swe was 1972 and 1974 and minimum swe was in 1987, 1992, 1994, and 2001. Average is for the period 1971 through 2000.

Mountain precipitation during February was 96 percent of average and 89 percent of last year. Valley precipitation during February was 82 percent of average and 288 percent of last year. Mountain and valley water year precipitation, beginning October 1, 2006, was 103 percent of average and 91 percent of last year.

Bitterroot Precipitation



Painted Rocks Lake storage was 130 percent of average and 134 percent of last year and Como storage was 189 percent of average and 221 percent of last year.

Assuming average precipitation March through July, April through July streamflows are forecast to average 80 percent to 88 percent.

Surface Water Supply Index (SWSI) was -1.4 in the Bitterroot River.

BITTERROOT RIVER BASIN
Streamflow Forecasts - March 1, 2007

Forecast Point	Forecast Period	<----- Drier ----- Future Conditions ----- Wetter ----->						30-Yr Avg. (1000AF)
		Chance Of Exceeding *						
		90% (1000AF)	70% (1000AF)	50% (1000AF)	(% AVG.)	30% (1000AF)	10% (1000AF)	
WF BITTERROOT nr Conner (2)	APR-JUL	84	103	117	82	132	156	143
	APR-SEP	92	113	128	82	144	170	157
BITTERROOT nr Darby	APR-JUL	280	340	385	84	430	505	460
	APR-SEP	330	390	430	84	475	545	515
COMO RESERVOIR Inflow	APR-JUL	63	70	75	96	80	88	78
	APR-SEP	66	73	78	95	83	90	82
BITTERROOT at Missoula	APR-JUL	880	1020	1130	90	1240	1410	1250
	APR-SEP	965	1120	1230	90	1350	1530	1370

BITTERROOT RIVER BASIN Reservoir Storage (1000 AF) - End of February					BITTERROOT RIVER BASIN Watershed Snowpack Analysis - March 1, 2007			
Reservoir	Usable Capacity	*** Usable Storage ***			Watershed	Number of Data Sites	This Year as % of	
		This Year	Last Year	Avg			Last Yr	Average
PAINTED ROCKS LAKE	31.7	9.1	6.8	7.0	WEST FORK BITTERROOT	3	72	75
COMO	align="right">34.9	align="right">23.4	align="right">10.6	align="right">12.4	EAST SIDE BITTERROOT	5	85	86
					WEST SIDE BITTERROOT	3	76	89
					BITTERROOT RIVER BASIN	10	79	85

* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.


The average is computed for the 1971-2000 base period.

- (1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
- (2) - The value is natural volume - actual volume may be affected by upstream water management.
- (3) - Median value used in place of average. The value listed under 30% is actually a 25% exceedance level. The value listed under 70% is actually a 75% exceedance level. Forecast issued in cooperation with Alberta Environment.

Lower Clark Fork River Basin

Snowpack conditions in the Lower Clark Fork River Basin were below average and increased 5 percent from February 1. Snow water content was 93 percent of average and 86 percent of last year. February accumulated snow water equivalent, as a percent of the seasonal peak, was 100 percent of average.


Lower Clark Fork Snow Water Equivalent



January maximum swe was established in 1997 and minimum swe was in 1977; February maximum swe was in 1972 and minimum swe was in 1977; March maximum swe was in 1972 and minimum was in 2005; April maximum swe was in 1972 and minimum swe was in 2005; May maximum swe was in 1972 and minimum swe was in 1977; and June maximum swe was in 1974 and minimum swe was in 1977. Average is for the period 1971 through 2000.

Mountain precipitation during February was 93 percent of average and 93 percent of last year. Valley precipitation during February was 110 percent of average and 159 percent of last year. Mountain and valley water year precipitation, beginning October 1, 2006, was 107 percent of average and 98 percent of last year.

Lower Clark Fork Precipitation



Noxon Rapids storage was 109 percent of average and 105 percent of last year.

Assuming average precipitation March through July, April through July streamflows are forecast to average 84 percent to 93 percent.

Surface Water Supply Index (SWSI) was -1.4 in the Clark Fork River below Bitterroot River and -0.1 in the Clark Fork River below Flathead River.

LOWER CLARK FORK RIVER BASIN
Streamflow Forecasts - March 1, 2007

		<<----- Drier ----- Future Conditions ----- Wetter ----->>						
Forecast Point	Forecast Period	Chance Of Exceeding *						30-Yr Avg. (1000AF)
		90% (1000AF)	70% (1000AF)	50% (1000AF)	(% AVG.)	30% (1000AF)	10% (1000AF)	
CLARK FORK blw Missoula	APR-JUL	1870	2200	2430	91	2660	2990	2660
	APR-SEP	2090	2450	2700	91	2950	3310	2960
CLARK FORK at St. Regis (1)	APR-JUL	2000	2870	3260	93	3650	4520	3520
	APR-SEP	2220	3180	3620	93	4060	5020	3910
CLARK FORK nr Plains (1,2)	APR-JUL	7070	8780	9560	95	10300	12100	10100
	APR-SEP	7760	9640	10500	95	11400	13200	11100
THOMPSON nr Thompson Falls	APR-JUL	139	170	193	94	215	255	205
	APR-SEP	162	195	220	96	245	285	230
PROSPECT CREEK at Thompson Falls	APR-JUL	82	96	107	92	118	135	116
	APR-SEP	89	104	115	93	126	144	124
CLARK FK at Whitehorse Rpds (1,2)	APR-JUL	7840	9810	10700	95	11600	13600	11300
	APR-SEP	8650	10800	11800	94	12800	15000	12500

LOWER CLARK FORK RIVER BASIN
Reservoir Storage (1000 AF) - End of February

LOWER CLARK FORK RIVER BASIN
Watershed Snowpack Analysis - March 1, 2007

Reservoir	Usable Capacity	*** Usable Storage ***			Watershed	Number of Data Sites	This Year as % of	
		This Year	Last Year	Avg			Last Yr	Average
NOXON RAPIDS	335.0	325.5	309.5	297.5	LOWER CLARK FORK BASIN	12	86	95


* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average is computed for the 1971-2000 base period.

- (1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
- (2) - The value is natural volume - actual volume may be affected by upstream water management.
- (3) - Median value used in place of average. The value listed under 30% is actually a 25% exceedance level. The value listed under 70% is actually a 75% exceedance level. Forecast issued in cooperation with Alberta Environment.


Jefferson River Basin

Snowpack conditions in the Jefferson River Basin were below average and increased 9 percent from February 1. Snow water content was 83 percent of average and 81 percent of last year. February accumulated snow water equivalent, as a percent of the seasonal peak, was 119 percent of average. Snowpack in the Ruby is tied for eighth lowest of record with 1964, 1973, and 1981, but has improved from fourth lowest of record on February 1.



January maximum swe was established in 1997 and minimum swe was in 1977; February maximum swe was in 1997 and minimum was in 1977; March maximum swe was in 1972 and minimum was in 1977; April maximum swe was in 1997 and minimum was in 1977; May maximum swe was in 1975 and minimum swe was in 1977; and June maximum swe was in 1982 and minimum in 1987. Average is for the period 1971 through 2000.

Mountain precipitation during February was 119 percent of average and 140 percent of last year. Valley precipitation during February was 204 percent of average and 271 percent of last year. Mountain and valley water year precipitation, beginning October 1, 2006, was 104 percent of average and 90 percent of last year.



Lima storage was 129 percent of average and 119 percent of last year; Clark Canyon storage was 80 percent of average and 118 percent of last year; and Ruby River storage was 96 percent of average and 123 percent of last year.

Assuming average precipitation March through July, April through July streamflows are forecast to average 56 percent to 76 percent.

Surface Water Supply Index (SWSI) was -2.0 in the Beaverhead River; -3.6 in the Ruby River; -0.8 in the Big Hole River; -1.0 in the Boulder River; and -0.2 in the Jefferson River as a whole.

JEFFERSON RIVER BASIN
Streamflow Forecasts - March 1, 2007

Forecast Point	Forecast Period	<<----- Drier ----- Future Conditions ----- Wetter ----->>						30-Yr Avg. (1000AF)
		Chance Of Exceeding *						
		90% (1000AF)	70% (1000AF)	50% (1000AF)	(% AVG.)	30% (1000AF)	10% (1000AF)	
LIMA RESERVOIR Inflow (2)	APR-JUL	38	53	66	69	82	114	96
	APR-SEP	38	54	69	66	88	124	104
CLARK CANYON Inflow (2)	APR-JUL	28	49	72	55	105	183	131
	APR-SEP	41	68	95	61	133	220	156
BEAVERHEAD RIVER at Barretts (2)	APR-JUL	0.0	52	99	59	146	215	168
	APR-SEP	0.0	73	129	65	185	270	200
RUBY RIVER Reservoir Inflow	APR-JUL	22	45	61	73	77	100	84
	APR-SEP	29	55	73	72	91	117	101
BIG HOLE RIVER at Wisdom	APR-JUL	28	74	104	86	134	178	121
	APR-SEP	29	78	111	85	144	192	130
BIG HOLE RIVER nr Melrose	APR-JUL	315	445	535	88	625	755	610
	APR-SEP	345	490	585	89	680	825	660
JEFFERSON nr Twin Bridges (2)	APR-JUL	260	475	620	79	765	980	785
	APR-SEP	330	545	695	79	845	1060	880
BOULDER RIVER nr Boulder	APR-JUL	40	58	72	92	89	118	78
	APR-SEP	43	62	78	92	97	128	85
WILLOW CREEK Reservoir Inflow	APR-JUL	1.8	9.0	13.8	77	18.4	25	17.9
	APR-SEP	1.5	9.8	15.4	77	21	29	20
JEFFERSON RIVER nr Three Forks (2)	APR-JUL	325	520	655	84	790	985	780
	APR-SEP	355	565	705	82	845	1050	860

JEFFERSON RIVER BASIN Reservoir Storage (1000 AF) - End of February					JEFFERSON RIVER BASIN Watershed Snowpack Analysis - March 1, 2007			
Reservoir	Usable Capacity	*** Usable Storage ***			Watershed	Number of Data Sites	This Year as % of	
		This Year	Last Year	Avg			Last Yr	Average
LIMA	84.0	46.0	38.8	35.7	BEAVERHEAD	16	83	84
CLARK CANYON	255.6	114.7	96.9	144.2	RUBY	11	81	79
RUBY RIVER	38.8	26.4	21.4	27.4	BIGHOLE	15	82	85
					BOULDER	9	79	84
					JEFFERSON RIVER BASIN	43	81	83


* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average is computed for the 1971-2000 base period.

- (1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
- (2) - The value is natural volume - actual volume may be affected by upstream water management.
- (3) - Median value used in place of average. The value listed under 30% is actually a 25% exceedance level. The value listed under 70% is actually a 75% exceedance level. Forecast issued in cooperation with Alberta Environment.


Madison River Basin

Snowpack conditions in the Madison River Basin were below average and increased 11 percent from February 1. Snow water content was 79 percent of average and 74 percent of last year and during February improved from fourth lowest of record to tenth lowest of record. February accumulated snow water equivalent, as a percent of the seasonal peak, was 118 percent of average.



January maximum swe was established in 1997 and minimum swe was in 1977; February maximum swe was in 1997 and minimum was in 1977; March maximum swe was in 1997 and minimum was in 1977; April maximum swe was in 1997 and minimum was in 1977; May maximum swe was in 1997 and minimum swe was in 1977; and June maximum swe was in 1995 and minimum in 2001. Average is for the period 1971 through 2000.

Mountain and valley precipitation during February was 122 percent of average and 147 percent of last year. Mountain and valley water year precipitation, beginning October 1, 2006, was 94 percent of average and 77 percent of last year.



Ennis Lake storage was 90 percent of average and 92 percent of last year and Hebgen Lake storage was 106 percent of average and 100 percent of last year.

Assuming average precipitation March through July, April through July streamflows are forecast to average 77 percent to 85 percent.

Surface Water Supply Index (SWSI) was -1.2 for the Madison River.

MADISON RIVER BASIN
Streamflow Forecasts - March 1, 2007

		<<----- Drier ----- Future Conditions ----- Wetter ----->>						
Forecast Point	Forecast Period	----- Chance Of Exceeding * -----						30-Yr Avg. (1000AF)
		90% (1000AF)	70% (1000AF)	50% (1000AF) (% AVG.)		30% (1000AF)	10% (1000AF)	

HEBGEN Reservoir Inflow	APR-JUL	260	305	335	86	365	410	390
	APR-SEP	345	395	430	86	465	515	500

ENNIS Reservoir Inflow (2)	APR-JUL	440	515	565	83	615	690	680
	APR-SEP	555	645	705	83	765	855	850

MADISON RIVER BASIN Reservoir Storage (1000 AF) - End of February					MADISON RIVER BASIN Watershed Snowpack Analysis - March 1, 2007			
Reservoir	Usable Capacity	*** Usable Storage ***			Watershed	Number of Data Sites	This Year as % of	
		This Year	Last Year	Avg			Last Yr	Average
ENNIS LAKE	41.0	28.2	30.7	31.4	MADISON abv HEBGEN LAKE	6	78	83
HEBGEN LAKE	377.5	281.8	280.5	265.2	MADISON blw HEBGEN LAKE	12	72	76
					MADISON RIVER BASIN	18	74	79

* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.


The average is computed for the 1971-2000 base period.

- (1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
- (2) - The value is natural volume - actual volume may be affected by upstream water management.
- (3) - Median value used in place of average. The value listed under 30% is actually a 25% exceedance level.
The value listed under 70% is actually a 75% exceedance level. Forecast issued in cooperation with Alberta Environment.

Gallatin River Basin

Snowpack conditions in the Gallatin River Basin were below average and increased 16 percent from February 1. Snow water content was 80 percent of average and 72 percent of last year ranking sixth lowest of record with 1994 but has improved from third lowest of record on February 1. February accumulated snow water equivalent, as a percent of the seasonal peak, was 147 percent of average.


Gallatin Snow Water Equivalent



January maximum swe was established in 1997 and minimum swe was in 1966; February maximum swe was in 1997 and minimum was in 1977; March maximum swe was in 1997 and minimum was in 1977, 1987 and 2005; April maximum swe was in 1997 and minimum was in 2001; May maximum swe was in 1970 and minimum swe was in 1987; and June maximum swe was in 1975 and minimum in 2001. Average is for the period 1971 through 2000.

Mountain precipitation during February was 137 percent of average and 156 percent of last year. Valley precipitation during February was 227 percent of average. Mountain and valley water year precipitation, beginning October 1, 2006, was 95 percent of average and 79 percent of last year.

Gallatin Precipitation



Middle Creek storage was 153 percent of average and 103 percent of last year.

Assuming average precipitation March through July, April through July streamflows are forecast to average 74 percent to 84 percent.

Surface Water Supply Index (SWSI) was -1.6 for the Gallatin River.

GALLATIN RIVER BASIN
Streamflow Forecasts - March 1, 2007

		<<----- Drier ----- Future Conditions ----- Wetter ----->>						
Forecast Point	Forecast Period	Chance Of Exceeding *						30-Yr Avg. (1000AF)
		90% (1000AF)	70% (1000AF)	50% (1000AF)	(% AVG.)	30% (1000AF)	10% (1000AF)	
GALLATIN RIVER nr Gateway	APR-JUL	280	335	370	84	405	460	440
	APR-SEP	320	385	425	83	465	530	515
HYALITE RESERVOIR Inflow	APR-JUL	14.9	17.9	20	91	22	25	22
	APR-SEP	17.6	21	23	92	25	28	25
GALLATIN RIVER at Logan (2)	APR-JUL	210	315	385	78	455	560	495
	APR-SEP	250	365	445	78	525	640	570

GALLATIN RIVER BASIN
Reservoir Storage (1000 AF) - End of February

Reservoir	Usable Capacity	*** Usable Storage ***		
		This Year	Last Year	Avg
MIDDLE CREEK	10.2	6.9	6.7	4.5

GALLATIN RIVER BASIN
Watershed Snowpack Analysis - March 1, 2007

Watershed	Number of Data Sites	This Year as % of	
		Last Yr	Average
UPPER GALLATIN	7	65	76
HYALITE	4	90	93
BRIDGER	2	66	68
GALLATIN RIVER BASIN	13	71	79


* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average is computed for the 1971-2000 base period.

- (1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
- (2) - The value is natural volume - actual volume may be affected by upstream water management.
- (3) - Median value used in place of average. The value listed under 30% is actually a 25% exceedance level. The value listed under 70% is actually a 75% exceedance level. Forecast issued in cooperation with Alberta Environment.


Missouri Mainstem River Basin

Snowpack conditions in the Headwaters Missouri Mainstem River Basin were below average and increased 11 percent from February 1. Snow water content was 93 percent of average and 89 percent of last year. February accumulated snow water equivalent, as a percent of the seasonal peak, was 131 percent of average.



January maximum swe was established in 1997 and minimum swe in 1977; February maximum swe was in 1972 and minimum swe was in 1977; March maximum swe in 1972 and minimum swe was in 1977; April maximum swe was in 1972 and minimum swe was in 1961, 1966, and 2005; May maximum swe was in 1975 and minimum swe was in 1977; and June maximum swe was in 1975 and minimum swe was in 1992. Average is for the period 1971 through 2000.

Mountain precipitation during February was 129 percent of average and 133 percent of last year. Valley precipitation during February was 212 percent of average and 256 percent of last year. Mountain and valley water year precipitation, beginning October 1, 2006, was 106 percent of average and 96 percent of last year.



Canyon Ferry Lake storage was 99 percent of average and 113 percent of last year; Helena Valley storage was 173 percent of average and 100 percent of last year; Lake Helena storage was 81 percent of average and 102 percent of last year; Hauser & Helena storage was 117 percent of average and 101 percent of last year; Holter Lake storage was 105 percent of average and 100 percent of last year; and Fort Peck Lake storage was 59 percent of average and 96 percent of last year.

Assuming average precipitation March through July, April through July streamflows are forecast to average 67 percent to 80 percent.

Surface Water Supply Index (SWSI) was -1.1 in the Missouri River above Canyon Ferry; -1.6 in the Missouri River below Canyon Ferry; --- in the Missouri River above Fort Peck; and -3.7 in the Missouri River below Fort Peck.

MISSOURI MAINSTEM RIVER BASIN
Streamflow Forecasts - March 1, 2007

		<----- Drier ----- Future Conditions ----- Wetter ----->						
Forecast Point	Forecast Period	Chance Of Exceeding *						30-Yr Avg. (1000AF)
		90% (1000AF)	70% (1000AF)	50% (1000AF)	(% AVG.)	30% (1000AF)	10% (1000AF)	
MISSOURI RIVER at Toston (2)	APR-JUL	990	1390	1660	81	1930	2330	2050
	APR-SEP	1070	1530	1850	77	2170	2630	2390
DEARBORN nr Craig	APR-JUL	50	79	97	80	116	144	121
	APR-SEP	53	83	102	82	121	151	125
MISSOURI RIVER at Fort Benton (2)	APR-JUL	1441	2024	2420	81	2817	3397	2990
	APR-SEP	1721	2417	2890	81	3361	4061	3570
MISSOURI RIVER at Virgelle (2)	APR-JUL	1700	2373	2830	82	3285	3960	3450
	APR-SEP	1990	2782	3320	82	3856	4651	4060
MISSOURI RIVER nr Landusky (2)	APR-JUL	1632	2351	2840	77	3330	4050	3690
	APR-SEP	1935	2784	3360	77	3936	4781	4350
MISSOURI RIVER below Fort Peck (2)	APR-JUL	1695	2424	2920	78	3416	4146	3740
	APR-SEP	1972	2816	3390	78	3966	4811	4330
LAKE SAKAKAWEA Inflow (2)	APR-JUL	5260	7159	8450	87	9744	11644	9740
	APR-SEP	4732	6916	8400	75	9881	12071	11200

MISSOURI MAINSTEM RIVER BASIN
Reservoir Storage (1000 AF) - End of February

Reservoir	Usable Capacity	*** Usable Storage ***		
		This Year	Last Year	Avg
CANYON FERRY LAKE	2043.0	1501.0	1331.0	1509.4
HELENA VALLEY	9.2	6.9	6.9	4.0
LAKE HELENA	12.7	10.9	10.7	13.4
HAUSER & HELENA	74.6	74.0	73.1	63.0
HOLTER LAKE	81.9	81.1	80.8	77.6
FORT PECK LAKE	18910.0	8651.0	9054.0	14728.0

MISSOURI MAINSTEM RIVER BASIN
Watershed Snowpack Analysis - March 1, 2007

Watershed	Number of Data Sites	This Year as % of	
		Last Yr	Average
HEADWATERS MAINSTEM	9	89	93
SMITH-JUDITH-MUSSELSHELL	14	88	102
SUN-TETON-MARIAS	14	97	90
MAINSTEM ab FT PECK RES	36	93	94
MILK RIVER BASIN	3	101	115
MISSOURI MAINSTEM BASIN	38	93	96

* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.


The average is computed for the 1971-2000 base period.

- (1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
- (2) - The value is natural volume - actual volume may be affected by upstream water management.
- (3) - Median value used in place of average. The value listed under 30% is actually a 25% exceedance level. The value listed under 70% is actually a 75% exceedance level. Forecast issued in cooperation with Alberta Environment.

Smith-Judith-Musselshell River Basins

Snowpack conditions in the Smith-Judith-Musselshell River Basins were near average and increased 23 percent from February 1. February accumulated snow water equivalent, as a percent of the seasonal peak, was 171 percent of average. Snow water content in the Smith River Basin was 94 percent of average and 83 percent of last year; in the Judith River Basin was 111 percent of average and 91 percent of last year; and in the Musselshell Basin River was 83 percent of average and 90 percent of last year. The Musselshell ranked seventh lowest of record with 1983, 1994, and 1977 but has improved from a record low on February 1.


Smith-Judith-Musselshell Snow Water Equivalent



January maximum swe was established in 1997 and minimum swe in 1988; February maximum swe was in 1978 and minimum swe was in 1987; March maximum swe was in 1978 and minimum swe was in 1987 and 2005; April maximum swe was in 1970 and minimum swe was in 1992; and May maximum swe was in 1970 and minimum swe was in 1987; and June maximum swe was in 1967 and minimum swe was in 1987, 1992, 1994, and 2001. Average is for the period 1971 through 2000.

Mountain and valley precipitation during February in the Smith-Belts was 141 percent of average and 122 percent of last year; in the Judith was 146 percent of average and 114 percent of last year; and in the Musselshell was 183 percent of average and 245 percent of last year. Mountain and valley water year precipitation, beginning October 1, 2006, was 114 percent of average and 96 percent of last year.

Smith-Judith-Musselshell Precipitation



Smith River storage was 77 percent of average and 134 percent of last year; Ackley Lake was 69 percent of average and 92 percent of last year; Bair storage was 45 percent of average and 75 percent of last year; Martinsdale storage was 0 percent of average; and Deadman's Basin was 77 percent of average and 116 percent of last year.

Assuming average precipitation March through July, April through July streamflows are forecast to average 48 percent to 73 percent.

Surface Water Supply Index (SWSI) was +0.4 in the Smith River; -0.7 in the Upper Judith River; and -2.1 in the Musselshell River.

SMITH-JUDITH-MUSSELSHELL RIVER BASINS
Streamflow Forecasts - March 1, 2007

		<<----- Drier ----- Future Conditions ----- Wetter ----->>						
Forecast Point	Forecast Period	Chance Of Exceeding *						30-Yr Avg (1000AF)
		90% (1000AF)	70% (1000AF)	50% (1000AF)	(% AVG.)	30% (1000AF)	10% (1000AF)	
SHEEP CREEK nr White Sulphur Spgs.	APR-JUL	7.9	11.5	13.8	81	16.2	19.7	17.1
	APR-SEP	9.8	13.8	16.6	83	19.4	23	20
SMITH RIVER abv Eagle Creek	APR-JUL	37	80	106	80	133	174	133
	APR-SEP	43	91	120	81	150	197	149
NF MUSSELSHELL nr Delpine	APR-JUL	1.1	2.6	3.6	78	4.6	6.1	4.6
	APR-SEP	1.4	3.1	4.2	78	5.3	7.0	5.4
SF MUSSELSHELL abv Martinsdale	APR-JUL	4.2	24	38	73	52	72	52
	APR-SEP	5.1	26	41	73	56	77	56
MUSSELSHELL at Harlowton (2)	APR-JUL	0.0	29	51	66	72	104	77
	APR-SEP	0.0	29	52	64	75	108	81
MUSSELSHELL nr Roundup (2)	APR-JUL	0.0	16.8	57	58	98	157	99
	APR-SEP	0.0	18.0	60	59	102	163	102

SMITH-JUDITH-MUSSELSHELL RIVER BASINS
Reservoir Storage (1000 AF) - End of February

SMITH-JUDITH-MUSSELSHELL RIVER BASINS
Watershed Snowpack Analysis - March 1, 2007

Reservoir	Usable Capacity	*** Usable Storage ***			Watershed	Number of Data Sites	This Year as % of	
		This Year	Last Year	Avg			Last Yr	Average
SMITH RIVER	10.6	5.1	3.8	6.6	SMITH	8	83	94
ACKLEY LAKE	7.0	2.2	2.4	3.2	HIGHWOOD	0	148	0
BAIR	7.0	1.8	2.4	4.0	JUDITH	6	91	111
MARTINSDALE	23.1	0.0	8.8	10.2	MUSSELSHELL	5	90	83
DEADMAN'S BASIN	72.2	38.2	32.8	49.8	SMITH-JUDITH-MUSSELSHELL	14	88	102

* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.


The average is computed for the 1971-2000 base period.

- (1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
- (2) - The value is natural volume - actual volume may be affected by upstream water management.
- (3) - Median value used in place of average. The value listed under 30% is actually a 25% exceedance level. The value listed under 70% is actually a 75% exceedance level. Forecast issued in cooperation with Alberta Environment.

Sun-Teton-Marias River Basins

Snowpack conditions in the Sun-Teton-Marias River Basins were below average and the percentage did not change from February 1. February accumulated snow water equivalent, as a percent of the seasonal peak, was 100 percent of average. Snow water content in the Sun River Basin was 91 percent of average and 96 percent of last year; in the Teton River Basin was 90 percent of average and 108 percent of last year; and in the Marias River Basin was 88 percent of average and 98 percent of last year.


Sun-Teton-Marias Snow Water Equivalent



January maximum swe was established in 1997 and minimum swe was in 1988; February maximum swe was in 1972 and minimum swe was in 2001; March maximum swe was in 1972 and minimum swe was in 2005; April maximum swe was in 1972 and minimum swe was in 2005; May maximum swe was in 1972 and minimum swe was in 1977; and June maximum was in 1972 and minimum swe was in 1987, 1992, and 1998. Average is for the period 1971 through 2000.

Mountain and valley precipitation during February in the Sun was 96 percent of average and 74 percent of last year; in the Teton was 132 percent of average and 95 percent of last year; and in the Marias was 90 percent of average and 72 percent of last year. Mountain and valley water year precipitation, beginning October 1, 2006, was 103 percent of average and 108 percent of last year.

Sun-Teton-Marias Precipitation



Gibson storage was 96 percent of average and 346 percent of last year; Pishkun storage was 0 percent of average; Willow Creek storage was 126 percent of average and 108 percent of last year; Lower Two Medicine Lake storage was not available; Four Horns Lake storage was not available; Swift storage was 124 percent of average and 131 percent of last year; Lake Frances storage was 37 percent of average and 71 percent of last year; and Lake Elwell (Tiber) storage was 103 percent of average and 92 percent of last year.

Assuming average precipitation March through July, April through July streamflows are forecast to average 76 percent to 95 percent.

Surface Water Supply Index (SWSI) was -0.3 in the Sun River; 0.0 in the Teton River; -3.0 in the Birch/Dupuyer Creeks; and ---- in the Marias River above Tiber Reservoir; and ---- in the Marias River below Tiber Reservoir.

SUN-TETON-MARIAS RIVER BASINS
Streamflow Forecasts - March 1, 2007

Forecast Point	Forecast Period	<<----- Drier ----- Future Conditions ----- Wetter ----->>						30-Yr Avg. (1000AF)
		Chance Of Exceeding *						
		90% (1000AF)	70% (1000AF)	50% (1000AF)	(% AVG.)	30% (1000AF)	10% (1000AF)	
GIBSON Reservoir Inflow	APR-JUL	335	405	450	98	495	565	460
	APR-SEP	380	450	495	98	540	610	505
TWO MEDICINE RIVER nr Browning	APR-JUL	123	166	195	95	222	267	205
	APR-SEP	132	176	205	95	235	280	215
BADGER CREEK nr Browning (2)	APR-JUL	61	82	97	98	112	133	99
	APR-SEP	75	97	113	98	129	151	115
SWIFT RESERVOIR Inflow	APR-JUL	34	51	62	97	73	90	64
	APR-SEP	45	62	74	96	86	103	77
DUPUYER CREEK nr Valier	APR-JUL	3.5	8.1	12.2	87	17.2	26	14.0
	APR-SEP	4.2	9.4	14.0	89	19.5	29	15.7
CUT BANK CREEK nr Browning	APR-JUL	54	67	76	99	85	98	77
	APR-SEP	60	74	83	99	92	106	84
MARIAS RIVER nr Shelby (2)	APR-JUL	247	340	400	96	460	555	415
	APR-SEP	265	360	420	96	480	575	440
TETON nr Dutton	APR-JUL	13.3	30	49	96	68	111	51
	APR-SEP	18.3	38	57	97	82	131	59

SUN-TETON-MARIAS RIVER BASINS
Reservoir Storage (1000 AF) - End of February

Reservoir	Usable Capacity	*** Usable Storage ***		
		This Year	Last Year	Avg
GIBSON	99.1	47.7	13.8	49.7
PISHKUN	32.0	0.0	17.4	17.6
WILLOW CREEK	32.2	28.9	26.8	23.0
LOWER TWO MEDICINE LAKE		NO REPORT		
FOUR HORNS LAKE		NO REPORT		
SWIFT	30.0	20.5	15.6	16.5
LAKE FRANCES	112.0	24.8	34.7	66.8
LAKE ELWELL (TIBER)	1347.0	652.3	707.4	631.9

SUN-TETON-MARIAS RIVER BASINS
Watershed Snowpack Analysis - March 1, 2007

Watershed	Number of Data Sites	This Year as % of	
		Last Yr	Average
SUN	7	96	91
TETON	4	108	90
MARIAS	6	98	88
SUN-TETON-MARIAS	15	98	90


* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average is computed for the 1971-2000 base period.

- (1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
- (2) - The value is natural volume - actual volume may be affected by upstream water management.
- (3) - Median value used in place of average. The value listed under 30% is actually a 25% exceedance level. The value listed under 70% is actually a 75% exceedance level. Forecast issued in cooperation with Alberta Environment.


St. Mary and Milk River Basins

Snowpack conditions in the St. Mary and Milk River Basins were near average and increased 20 percent from February 1. St. Mary February accumulated snow water equivalent, as a percent of the seasonal peak, was 100 percent of average. Snow water content in the St. Mary River Basin was 87 percent of average and 85 percent of last year and increased 1 percent from February 1. The Bearpaw Mountains were 115 percent of average and 101 percent of last year and increased 42 percent from February 1. The Cypress Hills in Canada were 152 percent of average and 240 percent of last year and increased 48 percent from February 1.





Bearpaw - January maximum swe was established in 1978 and minimum swe was in 1981; February maximum swe was 1978 and minimum was in 1973; March maximum swe was 1978 and minimum swe was 2005; April maximum swe was in 1985 and minimum swe was in 1973, 1983, and 1986; May maximum swe was 1975 and the minimum, 0.0, has occurred in several years. Average is for the period 1971 through 2000.

St. Mary - January maximum swe was established in 1997 and minimum swe was in 1988; February maximum swe was in 1972 and minimum swe was in 2001; March maximum swe was in 1972 and minimum swe was in 2005; April maximum swe was in 1967 and 1972 and minimum swe was in 2005; May maximum swe was in 1997 and minimum swe was in 1977; and June maximum swe was in 1972 and minimum swe was 1992. Average is for the period 1971 through 2000.



Mountain and valley precipitation in the St. Mary River Basin during February was 83 percent of average and 64 percent of last year with the water year precipitation, beginning October 1, 2006, at 108 percent of average and 98 percent of last year. The Milk River Basin during February was 198 percent of average and 109 percent of last year with the water year precipitation, beginning October 1, 2006, at 127 percent of average and 104 percent of last year.



Lake Sherburne storage was 139 percent of average and 92 percent of last year; Fresno storage was 77 percent of average and 82 percent of last year; Beaver Creek storage was 122 percent of average and 80 percent of last year; and Nelson storage was 79 percent of average and 76 percent of last year.

Assuming average precipitation March through July, April through July streamflows are forecast to average 77 percent to 100 percent.

Surface Water Supply Index (SWSI) was +1.3 in the St. Mary River and -0.3 in the Milk River.

ST. MARY and MILK RIVER BASINS
Streamflow Forecasts - March 1, 2007

Forecast Point	Forecast Period	<<----- Drier ----- Future Conditions ----- Wetter ----->>						30-Yr Avg. (1000AF)
		Chance Of Exceeding *						
		90% (1000AF)	70% (1000AF)	50% (1000AF)	(% AVG.)	30% (1000AF)	10% (1000AF)	
LAKE SHERBURNE Inflow	APR-JUL	88	96	102	97	108	116	105
	APR-SEP	102	111	118	97	125	134	122
ST. MARY RIVER nr Babb (2)	APR-JUL	320	350	375	97	400	430	385
	APR-SEP	375	415	440	98	465	505	450
ST. MARY RIVER at US/CAN Border (2)	APR-JUL	360	405	440	101	475	520	435
	APR-SEP	430	485	520	101	555	610	515
MILK RIVER @ Western Crossing (3)	MAR-JUL	10.7	25	38	93	53	72	41
	MAR-SEP	11.6	27	42	98	58	79	43
MILK RIVER @ Milk River, AB (2,3)	MAR-JUL	18.2	41	63	104	88	120	61
	MAR-SEP	19.0	45	69	105	96	132	66
MILK RIVER @ Eastern Crossing(2,3)	MAR-JUL	22	52	80	96	111	151	83
	MAR-SEP	24	56	87	99	121	166	88
BEAVER CREEK near Havre	MAR-JUL	3.8	7.3	10.5	109	14.5	22	9.6

ST. MARY and MILK RIVER BASINS
Reservoir Storage (1000 AF) - End of February

ST. MARY and MILK RIVER BASINS
Watershed Snowpack Analysis - March 1, 2007

Reservoir	Usable Capacity	*** Usable Storage ***			Watershed	Number of Data Sites	This Year as % of	
		This Year	Last Year	Avg			Last Yr	Average
LAKE SHERBURNE	64.3	39.7	43.3	28.6	ST. MARY	3	85	87
FRESNO	127.0	40.0	48.8	52.0	BEARPAW MOUNTAINS	3	101	115
BEAVER CREEK	3.5	2.8	3.5	2.3	CYPRESS HILLS, CANADA	0	0	0
NELSON	66.8	26.0	34.0	33.1	MILK RIVER BASIN	3	101	115
					ST. MARY & MILK BASINS	6	88	92

* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.


The average is computed for the 1971-2000 base period.

- (1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
- (2) - The value is natural volume - actual volume may be affected by upstream water management.
- (3) - Median value used in place of average. The value listed under 30% is actually a 25% exceedance level. The value listed under 70% is actually a 75% exceedance level. Forecast issued in cooperation with Alberta Environment.

Upper Yellowstone River Basin

Snowpack conditions in the Upper Yellowstone River Basin were below average and increased 11 percent from February 1. Snow water content was 81 percent of average and 79 percent of last year. February accumulated snow water equivalent, as a percent of the seasonal peak, was 127 percent of average. Shields basin snowpack was fifth lowest of record but has improved from lowest of record on February 1. Red Lodge, Rock Creek basins snowpack are tied at fourth lowest of record with 1977 and 1966 and slipped from sixth lowest of record on February 1.


Upper Yellowstone Snow Water Equivalent



January maximum swe was established in 1997 and minimum swe was in 1988; February maximum swe was in 1997 and minimum swe was in 2001; March maximum swe was in 1997 and minimum swe was in 2001; April maximum swe was in 1997 and minimum swe was in 2001; May maximum swe was in 1997 and minimum swe was in 1987; and June maximum swe was 1996 and minimum swe was in 2001. Average is for the period 1971 through 2000.

Mountain precipitation during February was 120 percent of average and 139 percent of last year. Valley precipitation during February was 130 percent of average and 269 percent of last year. Mountain and valley water year precipitation, beginning October 1, 2006, was 100 percent of average and 91 percent of last year.

Upper Yellowstone Precipitation



Cooney storage was 99 percent of average and 82 percent of last year.

Assuming average precipitation March through July, April through July streamflows are forecast to average 76 percent to 87 percent.

Surface Water Supply Index (SWSI) was 0.0 in the Yellowstone River above Livingston; -2.2 in the Shields River; -2.3 in the Boulder River; -2.2 in the Stillwater River; -3.1 in the Rock/Red lodge Creeks; -2.1 in the Clarks Fork River; and -0.9 in the Yellowstone River above Bighorn River.

UPPER YELLOWSTONE RIVER BASIN
Streamflow Forecasts - March 1, 2007

		<<----- Drier ----- Future Conditions ----- Wetter ----->>						
Forecast Point	Forecast Period	Chance Of Exceeding *						30-Yr Avg. (1000AF)
		90% (1000AF)	70% (1000AF)	50% (1000AF)	(% AVG.)	30% (1000AF)	10% (1000AF)	

YELLOWSTONE at Lake Outlet	APR-JUL	375	460	515	87	570	655	590
	APR-SEP	505	615	690	86	765	875	805
YELLOWSTONE RIVER at Corwin Springs	APR-JUL	1250	1410	1530	93	1650	1810	1650
	APR-SEP	1510	1710	1840	93	1970	2170	1970
YELLOWSTONE RIVER near Livingston	APR-JUL	1530	1670	1770	93	1870	2010	1900
	APR-SEP	1820	1990	2100	92	2210	2380	2280
SHIELDS RIVER nr Livingston	APR-JUL	35	77	105	72	133	175	145
	APR-SEP	35	83	116	72	149	195	162
BOULDER RIVER at Big Timber	APR-JUL	180	230	260	91	290	340	285
	APR-SEP	195	245	280	89	315	365	315
MYSTIC LAKE Reservoir Inflow (2)	APR-JUL	43	50	54	92	58	65	59
	APR-SEP	57	64	69	91	74	81	76
STILLWATER RIVER nr Absarokee (2)	APR-JUL	320	385	430	87	475	540	495
	APR-SEP	390	460	510	87	560	630	585
CLARKS FORK RIVER nr Belfry	APR-JUL	380	435	470	87	505	560	540
	APR-SEP	410	465	505	85	545	600	595
COONEY RESERVOIR Inflow (2)	APR-JUL	7.9	24	35	75	46	62	47
	APR-SEP	15.0	32	43	75	54	71	57
YELLOWSTONE RIVER at Billings (2)	APR-JUL	2170	2720	3090	88	3460	4010	3510
	APR-SEP	2550	3190	3630	88	4070	4710	4120

UPPER YELLOWSTONE RIVER BASIN
Reservoir Storage (1000 AF) - End of February

Reservoir	Usable Capacity	*** Usable Storage ***		
		This Year	Last Year	Avg
MYSTIC LAKE	21.0	3.8	3.8	4.0
COONEY	27.4	16.2	19.8	16.4

UPPER YELLOWSTONE RIVER BASIN
Watershed Snowpack Analysis - March 1, 2007

Watershed	Number of Data Sites	This Year as % of	
		Last Yr	Average
YELLOWSTONE ab LIVINGSTON	18	80	84
SHIELDS	4	68	72
BOULDER-STILLWATER	4	74	85
RED LODGE-ROCK CREEK	7	90	70
CLARK'S FORK	7	82	82
UPPER YELLOWSTONE BASIN	36	79	81


* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average is computed for the 1971-2000 base period.

- (1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
- (2) - The value is natural volume - actual volume may be affected by upstream water management.
- (3) - Median value used in place of average. The value listed under 30% is actually a 25% exceedance level. The value listed under 70% is actually a 75% exceedance level. Forecast issued in cooperation with Alberta Environment.


Lower Yellowstone River Basin

Snowpack conditions in the Lower Yellowstone River Basin were below average and rank sixth lowest of record with 1992 and 1993, but have improved from fourth lowest of record on February 1. Snow water content was 77 percent of average and 92 percent of last year and increased 7 percent from February 1. February accumulated snow water equivalent, as a percent of the seasonal peak, was 107 percent of average.



January maximum swe was established in 1997 and minimum swe was in 2000; February maximum swe was in 1997 and minimum swe was in 2001; March maximum swe was in 1986 and minimum swe was in 2001; April maximum swe was in 1986 and minimum swe was in 2001; May maximum swe was in 1997 and minimum swe was in 1981; and June maximum swe was in 1995 and minimum swe was in 2001. Average is for the period 1971 through 2000.

Mountain and valley precipitation during February was 121 percent of average and 137 percent of last year. Mountain and valley water year precipitation, beginning October 1, 2006, was 95 percent of average and 89 percent of last year.



Bighorn Lake storage was 93 percent of average and 93 percent of last year and Tongue River storage was 209 percent of average and 121 percent of last year.

Assuming average precipitation March through July, April through July streamflows are forecast to average 60 percent to 74 percent.

Surface Water Supply Index (SWSI) was -1.3 in the Bighorn River below Bighorn Lake; -1.1 in the Little Bighorn River; -1.3 in the Yellowstone River below Bighorn River; -1.6 in the Tongue River; and -2.0 in the Powder River.

LOWER YELLOWSTONE RIVER BASIN
Streamflow Forecasts - March 1, 2007

		<----- Drier ----- Future Conditions ----- Wetter ----->						
Forecast Point	Forecast Period	Chance Of Exceeding *						30-Yr Avg. (1000AF)
		90% (1000AF)	70% (1000AF)	50% (1000AF)	(% AVG.)	30% (1000AF)	10% (1000AF)	
BIGHORN RIVER nr St. Xavier (2)	APR-JUL	670	920	1090	68	1260	1510	1610
	APR-SEP	740	1015	1200	68	1390	1660	1760
LITTLE BIGHORN RIVER nr Hardin	APR-JUL	61	88	106	83	124	151	128
	APR-SEP	71	99	120	83	140	170	144
TONGUE RIVER nr Dayton (2)	APR-JUL	58	75	86	90	97	114	96
	APR-SEP	68	86	98	90	110	128	109
BIG GOOSE CREEK nr Sheridan	APR-JUL	19.0	33	42	81	51	65	52
	APR-SEP	27	41	50	83	59	73	60
LITTLE GOOSE CREEK nr Big Horn	APR-JUL	16.7	23	28	82	33	39	34
	APR-SEP	23	31	36	86	41	49	42
TONGUE RIVER RESERVOIR Inflow (2)	APR-JUL	86	142	180	82	218	273	220
	APR-SEP	100	159	200	80	240	300	250
YELLOWSTONE RIVER at Miles City (2)	APR-JUL	2940	3770	4340	81	4910	5740	5360
	APR-SEP	3420	4390	5050	81	5710	6680	6210
POWDER RIVER at Moorehead	APR-JUL	23	84	125	61	166	225	205
	APR-SEP	37	99	141	61	183	245	230
POWDER RIVER near Locate	APR-JUL	83	120	145	62	170	207	235
	APR-SEP	92	132	160	62	188	227	260
YELLOWSTONE RIVER nr Sidney (2)	APR-JUL	2830	3690	4270	78	4850	5710	5480
	APR-SEP	3250	4230	4900	78	5570	6550	6280

LOWER YELLOWSTONE RIVER BASIN
Reservoir Storage (1000 AF) - End of February

LOWER YELLOWSTONE RIVER BASIN
Watershed Snowpack Analysis - March 1, 2007

Reservoir	Usable Capacity	*** Usable Storage ***			Watershed	Number of Data Sites	This Year as % of	
		This Year	Last Year	Avg			Last Yr	Average
BIGHORN LAKE	1356.0	769.3	829.5	826.3	WIND RIVER (Wyoming)	20	87	71
TONGUE RIVER	79.1	51.5	42.4	24.6	SHOSHONE RIVER (Wyoming)	6	88	72
					BIGHORN RIVER (Wyoming)	20	93	79
					LITTLE BIGHORN (Wyoming)	3	116	94
					TONGUE RIVER (Wyoming)	10	108	94
					POWDER RIVER (Wyoming)	9	82	75
					LOWER YELLOWSTONE BASIN (	48	93	77

* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average is computed for the 1971-2000 base period.

- (1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
- (2) - The value is natural volume - actual volume may be affected by upstream water management.
- (3) - Median value used in place of average. The value listed under 30% is actually a 25% exceedance level. The value listed under 70% is actually a 75% exceedance level. Forecast issued in cooperation with Alberta Environment.



Federal Building, Room 443
10 E. Babcock
Bozeman, MT 59715



Montana
Water Supply Outlook
Report
Natural Resources Conservation Service
Bozeman, MT

